



DUAL SEQUENTIAL LAUNCH

May 14, 2019 @ 0530hrs

RESEARCH

ACP and PCP Dose-VF Research Medical Directive

An ACP/PCP Paramedic may provide the treatment prescribed in this medical directive if certified and authorized.

INDICATIONS

Refractory Ventricular Fibrillation

CONDITIONS


Dual Sequential Defibrillation	
Age	≥ 18 years
LOA	Altered
HR	N/A
RR	N/A
SBP	N/A
Other	Non-traumatic VF of presumed cardiac etiology Presenting rhythm of VF Three previous consecutive shocks by treating paramedics Currently randomized to Dual Sequential arm of trial

CONTRAINDICATIONS

Dual Sequential Defibrillation	
Traumatic cardiac arrest	
Suspected overdose	
Cardiac arrest secondary to environmental cause (drowning, hypothermia, hanging)	
Any verbal or written DNR	
Any non-VF rhythm or ROSC during first three analyses by participating paramedics	

TREATMENT

Consider Dual Sequential Defibrillation	
	Age ≥ 18 Years
Dose	1 Dual Sequential Shock
Max. Single Dose	200J per Defibrillator
Dosing Interval	2 min
Max. # of Doses	3



ePCR Procedure Code
933.4 - DOSE VF
Dual Sequential Defibrillation Arm